



MEETING ABSTRACT

Open Access

Economic crisis and smoking in health professionals in Greece

Eleni Litsiou^{1*}, Aikaterini Tsoutsas¹, Vasiliki Saltagianni¹, Dimos Fotopoulos², Stavroula Kolokytha¹, Spyridon Zakynthinos¹, Paraskevi Katsaounou¹

From 11th Annual Conference of the International Society for the Prevention of Tobacco Induced Diseases (ISPTID)

Athens, Greece. 9-11 December 2013

Background

Despite the reduction of consumer disposable income due to the economic crisis which negatively affected tobacco (Euromonitor, 2012), the proportion of smokers in Greece is still the highest among all European Countries (42.0%). This percentage is even high in medical students (35.0%) and professionals [1,2].

Materials and methods

In our study, we investigated how occupational factors and economic crisis affect smoking in health professionals in Evangelismos Hospital (the largest in Greece). Our questionnaires included smoking history, demographic factors, working data, intention to quit and 4DSQ (measuring depression anxiety and related psychosomatic symptoms). In our sample of 500 participants (men/women = 1:4, 37.0% medical doctors, 58.0% nurses), 49.0% were smokers. Initially, we explored work factors affected by crisis. The amount of work has increased for 45% of workers with a parallel decrease of 10.0-20.0% in the income of 90.0%. After collecting the questionnaires we distributed informative leaflets about our Smoking Cessation Outpatient Clinic (SCC).

Results

Taking into account that a smoker (1 pack/d) spends on average 10.0-20.0% of his salary for smoking, it is a paradox that only 25.0% of them report an intention to quit. This could be explained because smokers addicted to nicotine use smoking for handling stress and depression. Consequently, our SCC focused in informing the staff that nicotine is a stimulant agent and thus is

wrongly interpreted as relaxant. We initiated a cognitive intervention-motivational coaching program in order to stop this vicious circle. Namely to dissociate smoking used for craving and alleviation of withdrawal symptoms from real-life stress. After being informed, 30.0% of smokers intend attempting smoking cessation in our clinic.

Conclusions

Further information and intervention programs are necessary so that smokers are convinced that nicotine is a stimulant agent and thus is wrongly interpreted as relaxant.

Acknowledgements

Excellence 2012.

Authors' details

¹Pulmonary department-ICU, Evangelismos Hospital, Athens, 10676, Greece.

²OKANA, Addiction Unit, Rethymnon, 74100, Greece.

Published: 6 June 2014

References

1. Caleyachetty A, Lewis S, McNeill A, Leonardi-Bee J: **Struggling to make ends meet: exploring pathways to understand why smokers in financial difficulties are less likely to quit successfully.** *Eur J Public Health* 2012, **22**(1):41-8.
2. Vardavas CI, Bouloukaki I, Linardakis MK, Tzilepi P, Tzanakis N, Kafatos AG: **Smoke-free hospitals in Greece: Personnel perceptions, compliance and smoking habit.** *TobInduc Dis* 2009, **31**(5):1-8.

doi:10.1186/1617-9625-12-S1-A23

Cite this article as: Litsiou et al.: Economic crisis and smoking in health professionals in Greece. *Tobacco Induced Diseases* 2014 **12**(Suppl 1):A23.

* Correspondence: elitsiou@yahoo.gr

¹Pulmonary department-ICU, Evangelismos Hospital, Athens, 10676, Greece

Full list of author information is available at the end of the article